

# Isotropic dispersion, line shape, and remnant Fermi surface in one hole problem

Z. Y. Weng, D. N. Sheng, and C. S. Ting

Texas Center for Superconductivity and Department of Physics  
University of Houston, Houston, TX 77204-5506

It is shown that the isotropic dispersion relation, line shape of the spectral function, and remnant Fermi surface found in recent photoemission experiment in insulating  $\text{Ca}_2\text{CuO}_2\text{Cl}_2$  by Ronning *et al.* (Science, **282**, 2067 (1998)) can be consistently explained by a *direct* holon hopping process which fundamentally undermines the conventional self-consistent Born approximation approach to the  $t - J$  model.

74.20.Mn, 79.60-i, 71.27.+a

The problem of one hole moving in the antiferromagnetic (AF) spin background can give us some important clue about the finite doping. For many years, the self-consistent Born approximation (SCBA) method [1–4] based on the  $t - J$  model has provided a standard description of the one hole problem, consistent with exact numerical calculations [5] up to 32 sites [6], which predicts a *sharp* quasiparticle coherent peak in the spectral function with an *anisotropic* dispersion relation around the Fermi points  $\mathbf{k}_0 = (\pm\pi/2, \pm\pi/2)$ .

But the SCBA results are not consistent with angle-resolved photoemission spectroscopy (ARPES) measurements in the parent compounds of high- $T_c$  superconductors:  $\text{Sr}_2\text{CuO}_2\text{Cl}_2$  [7,8] and  $\text{Ca}_2\text{CuO}_2\text{Cl}_2$  [9], which indicate that while the “quasiparticle” peak is *not* sharp at all with an intrinsic broad energy structure, its dispersion is fairly *isotropic* around the Fermi point  $\mathbf{k}_0$ . A further challenge to theory comes from the recent high-quality measurement [9] revealing the so-called remnant Fermi surface (RFS) in the electron *relative* momentum distribution  $n_k^r$ , in contrast to the low-energy Fermi point structure.

To explain the isotropy of the dispersion relation, the second and third neighbor hopping  $t'$  and  $t''$  adding to the  $t - J$  model were suggested in literature [8,10,11] which could improve the agreement between the theory and experiment under some proper choice of parameters. But Laughlin [12] has proposed a different view in which the isotropic dispersion may come from a more intrinsic origin: the spinon spectrum. In this spin-charge separation picture, the size of the dispersion is also accounted for by the superexchange  $J$  without  $t'$  and  $t''$ , and the injected hole decaying into a spinon-holon pair can provide a natural broad structure in the spectral function. Such a spin-charge separation scenario is fundamentally different from the Landau quasiparticle picture in the SCBA approach.

The purpose of this paper is to reexamine the one hole issue based on the  $t - J$  model. We show that due to the existence of a *direct* holon hopping term missing in the SCBA scheme, the low-energy physics can be completely changed from the SCBA prediction, and

the isotropic “quasiparticle” dispersion, line shape of the spectral function, as well as the remnant Fermi surface structure observed in experiment can all be explained as the consequence of the spin-charge separation, which in some main aspect is consistent with the proposal made in Ref. [12].

We start with the slave-fermion representation  $c_{i\sigma} = f_i^\dagger b_{i\sigma}$  [2]. In the  $t - J$  model  $H_{t-J} = H_t + H_J$ ,  $H_J = -\frac{J}{2} \sum_{\langle ij \rangle} (\hat{\Delta}_{ij}^s)^\dagger \hat{\Delta}_{ij}^s$  with  $\hat{\Delta}_{ij}^s = \sum_{\sigma} \bar{b}_{i\sigma} \bar{b}_{j-\sigma}$ . [Here  $\bar{b}_{i\sigma} \equiv (-\sigma)^i b_{i\sigma}$ , with  $(-\sigma)^i$  explicitly keeping track of the Marshall sign.] The motion of a doped hole is governed by  $H_t = -t \sum_{\langle ij \rangle} \hat{B}_{ji} f_i^\dagger f_j + H.c.$  with  $\hat{B}_{ji} = \sum_{\sigma} \sigma \bar{b}_{j\sigma}^\dagger \bar{b}_{i\sigma}$ . Even though the Schwinger-boson mean-field theory (SBMFT) [13] works quite well at half-filling, characterized by the bosonic resonating-valence-bond (RVB) order parameter  $\langle \hat{\Delta}_{ij}^s \rangle$ , the one hole problem is highly nontrivial. The reason can be attributed to the fact that the hopping integral  $\langle \hat{B}_{ji} \rangle = 0$  in  $H_t$ . In the SCBA approach, the hole has to acquire its kinetic energy through the *dynamic fluctuations* of  $\hat{B}_{ji}$  as follows. To the leading order in a large-S (spin wave) expansion, one may write [2]

$$\hat{B}_{ji} \approx \hat{B}_{ji}^{L,SW} = b_0 \sum_{\sigma} \sigma [\bar{b}_{j\sigma}^\dagger + \bar{b}_{i\sigma}], \quad (1)$$

where  $b_0$  denotes the condensed part of the Schwinger boson field, corresponding to AF long-range order (AFLRO) [14]. Then a hole created by  $\bar{c}_{i\sigma} = f_i^\dagger b_0 (-\sigma)^i$  can propagate by an SCBA assisted hopping and behave just like a Landau quasiparticle with four Fermi points at  $\mathbf{k}_0$  [1–4].

However, an important process is missing in the above picture in the spin-1/2 case. We will show below that due to the RVB nature of the SBMFT description [13], a *direct* hopping term for the holon actually exists but is omitted in (1). For the sake of clarity, one may set the temperature  $T$  at  $0^+$  where  $b_0$  (and the AFLRO) is switched off such that (1) vanishes. Then one can rewrite  $\hat{B}_{ji}$  in the following form  $\hat{B}_{ji} = (B_{ji}^0) e^{i\hat{\phi}_{ji}}$  where  $B_{ji}^0 = \sum_{\sigma} \bar{b}_{j\sigma}^\dagger \bar{b}_{i\sigma}$  and  $\hat{\phi}_{ji} = \pm(\pi/2)[1 - \sigma_{ji}]$  [ $\sigma_{ji} = \pm 1$  depending on  $\uparrow$  or  $\downarrow$  spinon exchanged with the holon at

the given link  $(ij)$ . If  $\langle B_{ji}^0 \rangle = B^0 \neq 0$ , the following *bare* holon hopping term will emerge

$$H_h = -t_{eff} \sum_{\langle ij \rangle} e^{i\hat{\phi}_{ji}} f_i^\dagger f_j + H.c., \quad (2)$$

with  $t_{eff} \equiv tB^0$ . Here  $\langle \hat{B}_{ji} \rangle = 0$  is entirely due to  $\langle e^{i\hat{\phi}_{ji}} \rangle = 0$  but the correct measuring of the gauge phase strength should be the gauge-invariant quantity  $e^{i\sum_r \hat{\phi}_{ji}}$  for a closed path  $\Gamma$ . One may write  $e^{i\sum_r \hat{\phi}_{ji}} = (-1)^{N_\Gamma^\downarrow}$  with  $N_\Gamma^\downarrow$  denoting the total number of  $\downarrow$  spins (or  $\uparrow$  spins by symmetry) “exchanged” with the hole along the path  $\Gamma$ . Here  $(-1)^{N_\Gamma^\downarrow}$  is nothing but the phase string factor first discussed in Ref. [15] in a rigorous formulation.

The proof of  $B^0 \neq 0$  is straightforward. At half-filling, one has the Bogoliubov transformation

$$\bar{b}_{i\sigma} = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} \eta_{\mathbf{k}\sigma} e^{i\sigma \mathbf{k} \cdot \mathbf{r}_i} \left[ u_{\mathbf{k}} \gamma_{\mathbf{k}\sigma} - v_{\mathbf{k}} \gamma_{\mathbf{k}-\sigma}^\dagger \right] \quad (3)$$

in which  $\gamma_{\mathbf{k}\sigma}$  represents an SBMFT elementary excitation operator, with a gapless spectrum at  $T = 0^+$  [13]

$$E_{\mathbf{k}} = 2.32J\sqrt{1 - \xi_{\mathbf{k}}^2}, \quad (4)$$

and  $u_{\mathbf{k}}^2 - v_{\mathbf{k}}^2 = 1$ ,  $2u_{\mathbf{k}}v_{\mathbf{k}} = \xi_{\mathbf{k}}/\sqrt{1 - \xi_{\mathbf{k}}^2}$  where  $\xi_{\mathbf{k}} = (\cos k_x + \cos k_y)/2$ . It is crucial to recognize that the factor  $\eta_{\mathbf{k}\sigma}$  in (3) which satisfies  $\eta_{\sigma} = \eta_{-\sigma}^*$  and  $|\eta| = 1$  cannot be completely determined at the mean-field level. Specifically, the mean-field order parameter  $\langle \bar{b}_{i\sigma} \bar{b}_{j-\sigma} \rangle$  is independent of different choices of  $\eta_{\mathbf{k}\sigma}$ . Then one may exploit this symmetry for the one hole case. At each fixed hole position, there is a sub-Hilbert-space for spins which at the mean-field level is described by the Bogoliubov transformation (3). By choosing  $\eta_{\mathbf{k}\sigma} = [-\text{sgn}(\xi_{\mathbf{k}})]^{k_h}$  where  $k_h = 0$  if the hole is on the even sublattice and  $k_h = 1$  if the hole is on the odd sublattice site, it is easy to find that  $B^0 = \frac{2}{N} \sum_{\mathbf{k} \neq 0} |\xi_{\mathbf{k}}| v_{\mathbf{k}}^2 \approx 0.4$  in  $H_t$  where a hole changing sublattices is involved, while the mean-field result remains the same for the superexchange term  $H_J$ . Note that such a hole-position-dependent choice of the phase  $\eta_{\mathbf{k}\sigma}$  is perfectly legitimate as in the sub-Hilbert-space in which the hole position is given, the commutation relations of  $\bar{b}$ ,  $\bar{b}^\dagger$  remain unchanged.

The existence of a bare hopping term  $H_h$  in (2) will fundamentally change the SCBA result. Note that one may express  $c_{i\sigma} = \bar{c}_{i\sigma} + \tilde{c}_{i\sigma}$  with  $\tilde{c}_{i\sigma} = f_i^\dagger : b_{i\sigma} :$  with  $: b_{i\sigma} \equiv b_{i\sigma} - b_0(-\sigma)^i$ . Due to (2), the propagator for  $\tilde{c}_{i\sigma}$  has a leading contribution  $\tilde{G}_e \approx iG_f \cdot G_b$  which reflects the spin-charge separation, where  $G_f = -i\langle T_t(f_i^\dagger(t)f_j(0)) \rangle$  and  $G_b = -i\langle T_t(: b_{i\sigma}(t) :: b_{j\sigma}^\dagger(0) :) \rangle$ . On the other hand, the Hamiltonian corresponding to the linear spin-wave approximation (1) may be rewritten as  $H_h^{LSW} = -t \sum_{\langle ij \rangle \sigma} \left( \bar{c}_{i\sigma}^\dagger \tilde{c}_{j\sigma} + \tilde{c}_{i\sigma}^\dagger \bar{c}_{j\sigma} \right) + H.c.$ , which

literally shows a process for the quasiparticle  $\bar{c}_\sigma$  to *decay* into a holon-spinon pair represented by  $\tilde{c}_\sigma$ . When both holon and quasiparticle have no bare kinetic energy,  $H_h^{LSW}$  represents a virtual process and an SCBA treatment of it reproduces the well-known results [1–4]. But once the holon acquires a bare kinetic energy whereas the quasiparticle has none,  $H_h^{LSW}$  simply describes the break-up process of a quasiparticle, where the propagator  $\tilde{G}_e$  for  $\bar{c}_\sigma$  should become proportional to  $\tilde{G}_e$  to the leading order approximation. Note that any higher order corrections to the quasiparticle self-energy also add to the holon self-energy so that the initial instability is always maintained. Therefore, the SCBA approach is totally undermined once the spin-charge separation is present where the holon as a free particle has its bare kinetic energy.

The spin-charge separation reflected in  $\tilde{G}_e$  leads to the convolution law of the spectral function in the following form

$$A^e(\mathbf{k}, \omega) = \frac{1}{N} \sum_{\mathbf{k}' \neq 0} v_{\mathbf{k}'}^2 A^h(\mathbf{k}' - \mathbf{k}, -E_{\mathbf{k}'} - \omega), \quad (5)$$

where the r.h.s. is obtained after using the SBMFT result,  $\text{Im}G_b = -\pi[u_{\mathbf{k}}^2\delta(\omega - E_{\mathbf{k}}) - v_{\mathbf{k}}^2\delta(\omega + E_{\mathbf{k}})]$ , for the spinon part, while the holon spectral function  $A^h(\mathbf{k}, \omega) \neq 0$  at  $\omega \geq 0$  which is to be determined by  $H_h$  in (2). By noting that

$$e^{i\hat{\phi}_{ji}} = e^{i\mathbf{k}_0 \cdot (\mathbf{r}_j - \mathbf{r}_i)} e^{-i\delta\hat{\phi}_{ji}} \quad (6)$$

with  $\delta\hat{\phi}_{ji} = \pm(\pi/2)\sigma_{ji}$  and  $\langle \delta\hat{\phi}_{ji} \rangle = 0$ , one finds that  $A^h(\mathbf{k}, \omega)$  should be peaked at  $\mathbf{k} = \mathbf{k}_0$  at low energy, and correspondingly the spinon spectrum  $E_{\mathbf{k}+\mathbf{k}_0}$  will essentially determine the low-energy “edge” of the electron spectral function in (5). Here we note that  $\delta\hat{\phi}_{ji}$  will generally play a role causing the incoherent motion of the holon as opposed to the *coherent* spinon with a  $\delta$ -function-like spectral function. As an approximation, we first simply treat  $\delta\hat{\phi}_{ji}$  as a random flux field and determine  $A^h(\mathbf{k}, \omega)$  numerically. We note that the overall shape of  $A^e(\mathbf{k}, \omega)$  will *not* be sensitive to such an approximation as to be discussed below.

The typical behavior of  $A^e(\mathbf{k}, \omega)$  versus  $\omega$  is shown in Fig. 1 at different  $\mathbf{k}$ 's along  $(0, 0)$  to  $\mathbf{k}_0$ . The small arrows mark the energy of  $E_{\mathbf{k}+\mathbf{k}_0}$  which is indeed correlated with the position of the “coherent” peak or more precisely the edge of  $A^e(\mathbf{k}, \omega)$ . The line shape of the spectral function in Fig. 1 reflects essentially the *convolution* law of spinon and holon in a spin-charge separation context, which is strikingly similar to those found in the ARPES measurements [7–9]. In particular, the spinon spectrum  $E_{\mathbf{k}+\mathbf{k}_0}$  fits the ARPES data reasonably well over the whole Brillouin zone as shown in Fig. 2 along  $\mathbf{k}_0$  ( $\Sigma$ ) to  $(0, 0)$  ( $\Gamma$ ) and  $(0, \pi)$  ( $X$ ) (which are symmetric in our theory) with  $J = 0.14\text{eV}$ . A very flat (only about 13% change)  $E_{\mathbf{k}+\mathbf{k}_0}$  along  $(0, 0)$  to  $(0, \pi)$  is also in good agreement with the

experiment. Therefore, the theory can naturally account for the ARPES without  $t'$ ,  $t''$  terms.

In Fig. 1, the holon spectral function is computed at  $t_{eff} = 2J$  ( $t \approx 5J$ ) with the random (white noise) flux strength chosen between  $\pm 0.1\pi$  per plaquette in  $48 \times 48$  lattice. But we emphasize that the general line shape in Fig. 1 looks *insensitive* to those parameters which mainly affect  $A^h$ . This is because the broad feature of the spectral function is chiefly due to the convolution law which reflects the spin-charge separation [12], and the high energy part of  $A^e$  is essentially decided by the *bandwidth* ( $\sim 8t_{eff}$ ) of the holon spectrum, *not* the sharpness of  $A^h$  itself. Note that the coherent factor  $v^2$  and the “incoherence” of the holon caused by the gauge phase in (2) are crucial for the coherent spinon spectrum to “show up” in  $A^e(\mathbf{k}, \omega)$ . In the one-dimensional case, the phase string also plays a role to shape the one-hole spectral function as discussed in Ref. [16].

Beyond the random-flux treatment, the precise form of  $\delta\hat{\phi}_{ji}$  will be important in determining the *relative* change at different momentum of an energy-integrated  $A^e$ : the so-called relative momentum distribution [9] defined by  $n_k^r(\omega_0) = \int_{\omega_0}^0 \frac{d\omega}{\pi} A^e(k, \omega)$ . Here if  $\omega_0$  is taken to  $-\infty$ ,  $n_k^r$  will reduce to the momentum distribution  $n_k$  which should be featureless ( $= 1/2$ ) for the one-hole case [note that  $\langle c_{i\sigma}^\dagger c_{j\sigma} \rangle = \langle b_{i\sigma}^\dagger f_i f_j^\dagger b_{j\sigma} \rangle = \delta_{ij} \langle b_{i\sigma}^\dagger b_{i\sigma} \rangle$  using the non-double-occupancy constraint]. Thus, within the framework of the  $t - J$  model, the observed RFS feature in  $n_k^r$  [9] for a finite  $\omega_0$  ( $\sim 0.5eV$ ) has nothing to do with the half-filling ground-state and must solely come from the *dynamics* of the doped hole. We have already seen that the characteristic “Fermi-point”  $\mathbf{k}_0$  arises from the leading term of the gauge phase  $\hat{\phi}_{ji}$  in (6), and we will see that beyond the random flux approximation  $\delta\hat{\phi}_{ji}$  may be responsible for an additional momentum structure in  $n_k^r$ .

The gauge phase  $\hat{\phi}$  can be more precisely described as a nonlocal phase string effect as noted before. It may be clearly seen by formulating  $G_f$  according to Ref. [15] as follows

$$G_f(j, i; E) = \sum_{\{c\}\{\sigma_s\}} \langle \hat{S}_c(\{\sigma_s\}; E) \rangle (-1)^{N_c^\dagger}, \quad (7)$$

with  $\hat{S}_c \equiv \left( \prod_{s=1}^K G_J(t) \left[ \bar{b}_{m_{s-1}\sigma_s}^\dagger \bar{b}_{m_s\sigma_s} \right] \right) G_J$ ,  $G_J = 1/(E - H_J)$ , and  $N_c^\dagger = \sum_s (1 - \sigma_s)/2$ . In  $\hat{S}_c$ , the site  $m_s$  is at a path  $c$  that the holon travels, which connects sites  $i$  and  $j$ :  $m_0 = i, m_1, \dots, m_K = j$ . In particular, one can prove [15] that  $\langle \hat{S}_c(E) \rangle \geq 0$  at low energy such that the phase string effect introduced by  $(-1)^{N_c^\dagger}$  is basically *nonrepairable*. What we need to know is how such an oscillating phase string shapes the high energy part of  $A^e$  to determine  $n_k^r$ . Due to its nonrepairable nature, the *singular part* of the phase string effect can be explicitly built into the wave function through a unitary transformation [15], leading to  $c_{i\sigma} = a_{i\sigma} \times e^{i\Theta_i^{string}}$ . Here  $a_{i\sigma}$  refers to

the *elementary* charge and spin excitations whose details are not important to our problem below. The key thing is the phase shift  $\Theta_i^{string}$  which will then modulate the electron propagator by

$$e^{-i[\Theta_j^{string} - \Theta_i^{string}]} = e^{-i \int_c \cdot \hat{\mathbf{A}}^f(\mathbf{r})} \quad (8)$$

with  $\hat{\mathbf{A}}^f(\mathbf{r}) = \frac{1}{2} \sum_l [\sum_\sigma \sigma n_{l\sigma}^b - 1] \frac{\hat{\mathbf{z}} \times (\mathbf{r} - \mathbf{r}_l)}{|\mathbf{r} - \mathbf{r}_l|^2}$  in the equal-time limit. Such a phase string factor originated from  $\hat{\phi}_{ji}$  can be similarly evaluated using the method outlined in Ref. [17], which leads to four “incommensurate peaks” at  $(\pm\pi\kappa, 0)$  and  $(0, \pm\pi\kappa)$  in momentum space with  $\kappa \sim 1$  [18]. If this oscillating factor solely decides the momentum structure of  $n_k^r$  with the rest term in the propagator mainly contributing a broadening, then one gets a contour plot of  $n_k^r$  in Fig. 3 in terms of the superposition of aforementioned four peaks (with  $\kappa = 0.75$ ) at an arbitrary broadening (which is presumably controlled by the energy cutoff  $\omega_0$ : when  $\omega_0 \rightarrow -\infty$ , the broadening in the momentum space should also go to infinity such that  $n_{\mathbf{k}} = 1/2$ ). Note that the values of  $n_{\mathbf{k}}^r$  here are only for *relative* comparison. We see that the overall feature is in qualitative agreement with the experimental data [9] whose one quarter portion is also shown in the inset of Fig. 3 for comparison. Experimentally [9] the RFS is defined at the sharp drop of  $n_k^r$  in  $\mathbf{k}$ -space.

Finally, we point out that strictly speaking the AFLRO and RVB condensations only exist in the thermodynamic limit and both (1) and (2) do not directly apply to a finite system. In a small sample, both amplitude and phase fluctuations in  $\hat{B}_{ji}$  are expected to be important and cannot be distinguished like in (2) where  $B^0 \neq 0$ . And in this case a generalized SCBA scheme may be constructed to account for the leading order effect in the hole dynamics. This should be the reason why the small-size exact calculations have always identified a quasiparticle, consistent with the SCBA as well as power-method results [3,4,19]. However, such an SCBA quasiparticle is expected to eventually break up into holon-spinon composite at large lattice so long as there is a bosonic RVB condensation [13] in the thermodynamic limit to warrant (2).

In conclusion, a direct holon hopping term is identified in the one hole problem due to the RVB spin pairing in the SBMFT framework, which is omitted in the SCBA approach. Consequently a drastically different characterization of low-energy physics emerges: The SCBA quasiparticle becomes unstable and is replaced by a holon-spinon composite in which the holon is *incoherent* due to the phase string effect associated with the bare hopping, while the coherent *spinon* spectrum determines the *isotropic* “quasiparticle” dispersion relation. The line shape of the spectral function simply reflects the spin-charge decomposition, while the remnant Fermi surface is caused by the oscillating phase string effect at high

energy. These properties well explain the recent ARPES measurements within the framework of the  $t - J$  model.

### ACKNOWLEDGMENTS

We wish to acknowledge useful discussions with A. Ferraz, T. Tohyama, T.K. Lee, and especially C. Kim and F. Ronning who kindly provided their experimental data. We thank the hospitality of Centro Internacional de Fisica da Materia Condensada of Univ. de Brasilia and the Aspen Center for Physics where this work was initiated and finalized. The present work was supported in part by Texas ARP No.3652707, the Robert A. Welch foundation, and TCSUH.


Fig. 2. The quasiparticle spectra determined by the ARPES [9] (open square) and by the present theory (solid curve).

Fig. 3. The contour plot of the electron momentum distribution exhibiting a “remnant Fermi surface” structure. The experimental data [9] are shown in the insert at the upper right corner.

- 
- [1] S. Schmitt-Rink *et al.* Phys. Rev. Lett. **60**, 2793 (1988).
  - [2] C. L. Kane *et al.*, Phys. Rev. B **39**, 6880 (1989).
  - [3] G. Martinez and P. Horsch, *ibid.*, B **44**, 317 (1991).
  - [4] Z. Lui and E. Manousakis, *ibid.*, B **44**, 2414 (1991).
  - [5] see, E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994) and references therein.
  - [6] P.W. Leung and R.J. Gooding, Phys. Rev. B **52**, R15711 (1995).
  - [7] B.O. Wells *et al.*, Phys. Rev. Lett. **74**, 964 (1995).
  - [8] C. Kim *et al.*, Phys. Rev. Lett. **80**, 4245 (1998).
  - [9] F. Ronning *et al.* Science, **282**, 2067 (1998).
  - [10] A. Nazarenko *et al.*, Phys. Rev. B **51**, 8676 (1995); R. Eder *et al.*, Phys. Rev. B **55**, R3414 (1997).
  - [11] G. B. Martins *et al.*, Phys. Rev. B **60**, R3716 (1999); T. Tohyama *et al.*, cond-mat/9904231 and references therein.
  - [12] R.B. Laughlin, Phys. Rev. Lett. **79**, 1726 (1997); J. Phys. Chem. Solids. **56**, 1627 (1995).
  - [13] D.P. Arovas and A. Auerbach, Phys. Rev. B **38**, 316 (1988).
  - [14] For convenience, the magnetization here is to lie in the  $x - y$  plane instead of along the  $z$ -axis as in Ref. [2].
  - [15] Z. Y. Weng *et al.*, Phys. Rev. B **55**, 3894 (1997); D. N. Sheng *et al.*, Phys. Rev. Lett. **77**, 5102 (1996).
  - [16] H. Suzuura and N. Nagaosa, Phys. Rev. B **56**, 3548 (1997).
  - [17] Z.Y. Weng *et al.*, Phys. Rev. B **59**, 11367(1999).
  - [18] Readers are referred to Ref. [17] for the detail, where a similar phase-string factor appearing in the spin-spin correlation function is discussed which gives rise to incommensurate shifts by  $(\pm 2\pi\delta\bar{\kappa}, 0)$ , etc., from  $(\pi, \pi)$  with  $\delta$  denoting the doping concentration and  $\bar{\kappa} \sim 1$ .
  - [19] T.K. Lee and C.T. Shih, Phys. Rev. B **55**, 5983 (1997).

Fig. 1. The theoretical electron spectral function. The arrows mark the corresponding energy of the isotropic spinon spectrum  $E_{\mathbf{k}+\mathbf{k}_0}$ .

FIG. 1



**FIG. 2**




FIG. 3

